


DEM Modeling of Ballistic Gelatin for Low Energy Impacts

4th Aspherix and CFDEM Conference

20–21 Apr 2023, Linz, Austria



HC Grobbelaar DNJ Els CJ Coetzee
*Dept of Mech & Mechatronic Eng,
Stellenbosch University, South Africa*

Contents

- 1 Blocks
- 2 Colours
- 3 Lists
- 4 Math



Stellenbosch
UNIVERSITY
IYUNIVESITHI
UNIVERSITEIT

Ballistic Gel

Blocks
Colours
Lists
Math

Blocks



Stellenbosch
UNIVERSITY
iYUNIVESITHI
UNIVERSITEIT

Ballistic Gel

Blocks

Colours

Lists

Math

General block

A general block ...

Alert block

An alert block ...







Example block

An example ...

Theorem (Theorem block)

A theorem ...

STB Colours

	Color name	RGB
	stbMaroon	(97, 34, 59)
	stbGold	(183, 153, 98)
	stbGreen	(130, 204, 174)
	stbOrange	(220, 68, 5)
	stbWine	(166, 10, 61)
	stbSoil	(100, 51, 53)



Stellenbosch
UNIVERSITY
UNIVERSITH
UNIVERSITEIT

Ballistic
Gel

Blocks

Colours

Lists

Math

Itemize

- First item
- Second item
- ...

Enumerate

- ① First item
- ② Second item
- ③ ...

Description

First item ...
Second item ...

... ...

Residue Theorem

Let f be analytic in the region G except for the isolated singularities a_1, a_2, \dots, a_m . If γ is a closed rectifiable curve in G which does not pass through any of the points a_k and if $\gamma \approx 0$ in G then

$$\frac{1}{2\pi i} \int_{\gamma} f = \sum_{k=1}^m n(\gamma; a_k) \text{Res}(f; a_k).$$

Another nice theorem from complex analysis is

Maximum Modulus

Let G be a bounded open set in \mathbb{C} and suppose that f is a continuous function on G^- which is analytic in G . Then

$$\max\{|f(z)| : z \in G^-\} = \max\{|f(z)| : z \in \partial G\}.$$

Thank you

